



St. JOHN'S

RESIDENTIAL PUBLIC SCHOOL

Sonagopalpur, Sampatchak, Patna-7

AFFILIATED TO C.B.S.E

CHEMISTRY NOTES

METALS AND NONMETALS

SCHOOL CODE --- 65403,

AFF.NO---330407

STD...Xth

PREPARED BYDR. S.K. VERMA

Alloys

Alloys are homogeneous mixtures of metal with other metals or nonmetals in fixed ratio. Alloy formation enhances the desirable properties of the material, such as **hardness, tensile strength and resistance to corrosion.**

Examples of a few alloys:

Brass: copper and zinc

Bronze: copper and tin


Solder: lead and tin

Amalgam: mercury and other metal


Corrosion

Gradual deterioration of material usually a metal by the action of **moisture, air or chemicals** in the surrounding environment.


Rusting:



Corrosion of copper:



Corrosion of silver:



Prevention of Corrosion

1. Coating with paints or oil or grease:

Application of paint or oil or grease on metal surfaces keep out air and moisture.

2. Alloying:

Alloyed metal is more resistant to corrosion. Example: stainless steel.

3. Galvanization:

This is a process of coating molten zinc on iron articles. Zinc forms a protective layer and prevents corrosion.

4. Electroplating:

It is a method of coating one metal with another by the use of electric current. This method not only lends protection but also enhances the metallic appearance.

Example: silver plating, nickel plating.

5. Sacrificial protection:

Magnesium is more reactive than iron. When it is coated on the articles made of iron or steel, it acts as the cathode, undergoes reaction (sacrifice) instead of iron and protects the articles.

Physical Properties of Metals

- Hard and have a high tensile strength
- Solid at room temperature
- Sonorous
- Good conductors of heat and electricity
- Malleable, i.e., can be beaten into thin sheets
- Ductile, i.e., can be drawn into thin wires
- High melting and boiling points (except Caesium (Cs) and Gallium (Ga))
- Density, (except alkali metals). Osmium - highest density and lithium - least density
- Lustrous
- Silver-grey in colour, (except gold and copper)

Non-Metals

Nonmetals are those elements which do not exhibit the properties of metals.

Physical Properties of Non-metals

- Occur as solids, liquids and gases at room temperature
- Brittle
- Non-malleable
- Non-ductile
- Non-sonorous
- Bad conductors of heat and electricity

Exceptions in Physical Properties

- Alkali metals (Na, K, Li) can be cut using a knife.
- Mercury is a liquid metal.
- Lead and mercury are poor conductors of heat.
- Mercury expands significantly for the slightest change in temperature.
- Gallium and caesium have a very low melting point.
- Iodine is non-metal but it has lustre.
- Graphite conducts electricity.
- Diamond conducts heat and has a very high melting point.


Chemical Properties of Metals

- Alkali metals (Li, Na, K, etc) react vigorously with water and oxygen or air.
- Mg reacts with hot water.
- Al, Fe and Zn react with steam.
- Cu, Ag, Pt, Au do not react with water or dilute acids.



Reaction of Metals with Oxygen (Burnt in Air)

Metal + Oxygen → Metal oxide (basic)

- Na and K are kept immersed in kerosene oil as they react vigorously with air and catch fire.





- Mg, Al, Zn, Pb react slowly with air and form a protective layer that prevents corrosion.



- Silver, platinum and gold don't burn or react with air.

Basic Oxides of Metals


Some metallic oxides get dissolved in water and form alkalis. Their aqueous solution turns red litmus blue.






Amphoteric Oxides of Metals

Amphoteric oxides are metal oxides which react with both acids as well as bases to form salt and water.

For example - **Al₂O₃, ZnO, PbO, SnO**





Reactivity Series

The below table illustrates the reactivity of metals from high order to low order.

Symbol	Element
K	Potassium (Highly Active Metal)
Na	Sodium
Ca	Calcium
Ba	Barium
Mg	Magnesium
Al	Aluminium
Zn	Zinc
Fe	Iron
Ni	Nickel
Sn	Tin
Pb	Lead
H	Hydrogen
Cu	Copper
Hg	Mercury
Ag	Silver
Au	Gold
Pt	Platinum

TRICK TO REMEMBER REACTIVITY SERIES OF METALS

(ALPHABET IS NOT A SYMBOL OF METAL)

ALWAYS REMEMBER AN IMAGINARY NAME

MR.

P----- (POTASSIUM)

S----- (SODIUM)

C -----(CALSIUM)

B -----(BARIUM)

M----- (MAGNESIUM)

A -----(ALUMINIUM)

Z----- (ZINC)

I----- (IRON)

N -----(NICKEL)

T----- (TIN)

L----- (LEAD)

H -----(HYDROGEN)

Cu----- (COPPER)

Hg----- (MERURY)






Ag----- (SILVER)

Au----- (GOLD)

Pt----- (PLATINUM)



Reaction of Metals with Water or Steam

Metal+Water→Metalhydroxide or Metaloxide+Hydrogen





Reaction of Metals with Acid

Metal+diluteacid→Salt+Hydrogengas



Hydrogen gas is not evolved when a metal reacts with NITRIC ACID, because HNO_3 is a strong oxidizing agent. It oxidizes the H_2 produced to water and itself gets reduced to any of the NITROGEN OXIDES (N_2O , NO , NO_2).

Only Mg and Mn, react with very dilute nitric acid to liberate hydrogen gas





Displacement Reaction

A more reactive element displaces a less reactive element from its compound or solution.



How Do Metals React with Solution of Other Metal Salts ?

Metal A + Salt of metal B → Salt of metal A + Metal B




Reaction of Metals with Bases

Base + metal → salt + hydrogen



Uses of displacement reaction

1. Extraction of metals
2. Manufacturing of steel
3. **Thermit reaction:** (Reaction between Aluminium powder and Iron III oxide)



The thermit reaction is used in welding of railway tracks, cracked machine parts, etc. It is highly exothermic displacement reaction. The amount of heat evolved is so large that converts the metals into molten state like IRON.

Extraction of Metals (METALLURGY)

The process of extraction of Metals from its ORES is called METALLURGY.


Occurrence of Metals

Most of the elements, especially metals occur in nature in the combined state with other elements. All these compounds of metals are known as **minerals**.

Those **MINERALS** from which we obtained **METALS** profitably is called **ores**.

Au, Pt - exist in the native or free state.

Extraction of Metals




Metals of high reactivity - Na, K, Mg, Al.

Metals of medium reactivity - Fe, Zn, Pb, Sn.

Metals of low reactivity - Cu, Ag, Hg





Roasting

Converts sulphide ores into oxides on heating strongly in the presence of excess air. It also removes volatile impurities.



Calcination




Converts carbonate and hydrated ores into oxides on heating strongly in the presence of limited air. It also removes volatile impurities.






Extracting Metals Low in Reactivity Series

By self-reduction-


when the sulphide ores of less electropositive metals like Hg, Pb, Cu etc., are heated in air, a part of the ore gets converted to oxide which then reacts with the remaining sulphide ore to give the crude metal and sulphur dioxide. In this process, no external reducing agent is used.







Extracting Metals in the Middle of Reactivity Series

Smelting - it involves heating the roasted or calcined ore (metal oxide) to a high temperature with a suitable reducing agent. The crude metal is obtained in its molten state.



Aluminothermic reaction -

also known as the Goldschmidt reaction is a highly exothermic reaction in which metal oxides usually of Fe and Cr are heated to a high temperature with aluminium.




Extraction of Metals Towards the Top of the Reactivity Series

Electrolytic reduction:

1. Down's process: (NOT IN SYLLABUS)


1. Molten NaCl is electrolysed in a special apparatus.

At the **cathode** (reduction):



Metal is deposited.

At the **anode** (oxidation):




Chlorine gas is liberated.

2. Hall's process: (NOT IN SYLLABUS)


Mixture of molten alumina and a fluoride solvent usually cryolite, (Na_3AlF_6) is electrolysed.

At the **cathode** (reduction):



Metal is deposited.

At the **anode** (oxidation):



Oxygen gas is liberated.

Enrichment or Concentration of Ores

It means the removal of **impurities or gangue** from ore, through various physical and chemical processes. The technique used for a particular ore depends on the difference in the properties of the ore and the gangue.

Refining of Metals

Removing **impurities or gangue** from crude metal. It is the last step in metallurgy and is based on the difference between the properties of metal and the gangue.

Electrolytic Refining

Metals like copper, zinc, nickel, silver, tin, gold etc., are refined electrolytically.

Anode: impure or crude metal

Cathode: a thin strip of pure metal

Electrolyte: aqueous solution of metal salt

From anode (oxidation): metal ions are released into the solution

At cathode (reduction): the equivalent amount of metal from solution is deposited
Impurities deposit at the bottom of the anode.

Electronic Configuration

Group 1 elements – Alkali metals

Element	Electronic Configuration
Lithium(Li)	2,1
Sodium(Na)	2,8,1
Potassium(K)	2,8,8,1
Rubidium(Rb)	2,8,18,8,1

Group 2 elements – Alkaline earth metals

Element	Electronic Configuration
Beryllium(Be)	2,2
Magnesium(Mg)	2,8,2
Calcium(Ca)	2,8,8,2
Stronium(Sr)	2,8,18,8,2

How Do Metals and Nonmetals React


Metals lose valence electron(s) and form **cations**.

Non-metals gain those electrons in their valence shell and form **anions**.


The **cation** and the **anion** are attracted to each other by **strong electrostatic force**, thus forming an **ionic bond or Electrovalent Bond**.

For example: In calcium chloride, the ionic bond is formed by opposite charged calcium and chloride ions.

Calcium atom loses 2 electrons and attains the electronic configuration of the nearest noble gas (Ar). By doing so, it gains a net charge of +2.



The two Chlorine atoms take one electron each, thus gaining a charge of -1 (each) and attain the electronic configuration of the nearest noble gas (Ar).



Ionic Compounds or Electrovalent compounds

The electrostatic attractions between the opposite charged ions hold the compound together.

Example: MgCl_2 , CaO , MgO , NaCl etc.

Properties of Ionic Compound

Ionic compounds

1. Are usually crystalline solids (made of ions).
2. Have high melting and boiling points.
3. Conduct electricity when in aqueous solution and when melted.
4. Are mostly soluble in water and polar solvents.

Physical Nature


Ionic solids usually exist **in regular, well-defined crystal structures.**

Electric Conduction of Ionic Compounds

Ionic compounds conduct electricity in the molten or aqueous state when ions become free and act as charge carriers.

In solid form, ions are strongly held by electrostatic forces of attractions and are not free to move; hence do not conduct electricity.

In solid state



In molten and aqueous solution




● ions move to the negative terminal

● ions move to the positive terminal

For example,


ionic compounds such as NaCl does not conduct electricity when solid but when dissolved in water or in a molten state, it will conduct electricity.



Melting and Boiling Points of Ionic Compounds

In ionic compounds, the strong electrostatic forces between ions require a high amount of energy to break. Thus, the melting point and boiling point of an ionic compound are usually very high.

Solubility of Ionic Compounds



Most ionic compounds are soluble in water due to the separation of ions by water. This occurs due to the polar nature of water.

For example,

NaCl is a 3-D salt crystal composed of Na⁺ and Cl⁻ ions bound together through **electrostatic forces of attractions**. When a crystal of NaCl comes into contact with water, the partial positively charged ends of water molecules interact with the Cl⁻ ions, while the negatively charged end of the water molecules interacts with the Na⁺ ions.

This ion-dipole interaction between ions and water molecules assist in the breaking of the strong electrostatic forces of attractions within the crystal and ultimately in the solubility of the crystal.

.....

.....